

High Resolution CO and H₂ Molecular Line Imaging of a Cometary Globule in the Helix Nebula¹

P. J. Huggins², T. Forveille^{3,4}, R. Bachiller⁵, P. Cox⁶, N. Ageorges⁷, J. R. Walsh⁸

ABSTRACT

We report high resolution imaging of a prominent cometary globule in the Helix nebula in the CO $J = 1 - 0$ (2.6 mm) and H₂ $v = 1 - 0$ $S(1)$ (2.12 μm) lines. The observations confirm that globules consist of dense condensations of molecular gas embedded in the ionized nebula. The head of the globule is seen as a peak in the CO emission with an extremely narrow line width (0.5 km s⁻¹) and is outlined by a limb-brightened surface of H₂ emission facing the central star and lying within the photo-ionized halo. The emission from both molecular species extends into the tail region. The presence of this extended molecular emission provides new constraints on the structure of the tails, and on the origin and evolution of the globules.

Subject headings: Planetary nebulae: general – Planetary nebulae: individual: NGC 7293 – Stars: AGB and post-AGB

1. Introduction

The cometary globules in the Helix nebula (NGC 7293) are among the most remarkable structures seen in planetary nebulae (PNe). They occur in large numbers in the lower

¹Based on observations carried out with the IRAM Plateau de Bure Interferometer. IRAM is supported by INSU/CNRS (France), MPG (Germany) and IGN (Spain).

²Physics Department, New York University, 4 Washington Place, New York, NY 10003, USA

³CFHT, PO Box 1597, Kamuela, HI 96743, USA

⁴Observatoire de Grenoble, B.P. 53X, 38041 Grenoble Cedex, France

⁵IGN Observatorio Astronómico Nacional, Apartado 1143, E-28800 Alcalá de Henares, Spain

⁶Institut d'Astrophysique Spatiale, Université de Paris XI, 91405 Orsay, France

⁷European Southern Observatory, Alonso de Córdova 3107, Santiago, Chile

⁸Space Telescope Coordinating Facility, ESO, D-85748 Garching bei München, Germany

ionization regions of the nebula, and appear in high resolution optical images as small ($\sim 1''$), convex, photo-ionized surfaces facing the central star, with comet-like tails extending in the opposite direction (e.g., Meaburn et al. 1992; O’Dell & Handron 1996). These structures are seen in other PNe (e.g., O’Dell et al. 2002) and are probably quite common, but they are best seen in the Helix nebula because it is the nearest example, at a distance of $D \sim 200$ pc (parallax = 4.70 ± 0.75 mas, Harris et al. 1997).

A key step in understanding the nature of the Helix globules has been their detection in CO by Huggins et al. (1992). With a resolution $\sim 12''$, the CO observations did not resolve their structure, but demonstrated that globules contain a major mass component of molecular gas, consistent with observations of dust, seen in absorption against the nebula emission by Meaburn et al. (1992). The molecular gas places important constraints on the origin and evolution of the globules, and provides a direct connection with the massive shell of neutral gas that surrounds the ionized nebula (Forveille & Huggins 1991; Young et al. 1999; Rodriguez, Goss, & Williams 2002). This connection is underscored by wide field imaging of the nebula in the infrared lines of H_2 by Kastner et al. (1996), Cox et al. (1998), and Speck et al. (2002), at resolutions from $8''$ to $2''$, that show a highly fragmented envelope.

In order to determine the detailed relation between the molecular gas and the structure of the globules revealed by optical images, we have made new observations of the Helix nebula in both CO and H_2 with significantly better resolution than previous observations. In this *Letter*, we report results on the molecular gas in a single cometary globule that resolve its head-tail structure.

2. Observations

The globule observed is a prominent feature lying within the ionized nebula to the north of the central star at offsets ($-10''$, $+135''$). It is designated C1 by Huggins et al. (1992) and its location is shown in Fig. 3a of Meaburn et al. (1998) where it is labeled “1”. Optical images of the globule from Walsh & Meaburn (1993) in $H\alpha + [N\text{ II}] 6584 \text{ \AA}$ and in $[O\text{ III}] 5007 \text{ \AA}$, where it is seen in absorption against the nebular emission, are shown in the top right panels of Fig. 1.

The CO observations were made in the 2.6 mm (115 GHz) $J = 1 - 0$ line using the IRAM interferometer at Plateau de Bure, France, in May 1999. The array consisted of five, 15 m antennas, equipped with SIS heterodyne receivers. The observations were made with the D configuration of the array, with maximum baselines of ~ 147 m. The primary

beam size of the interferometer is $44''$ at 2.6 mm, and the effective velocity resolution for the analysis is 0.2 km s^{-1} . The uv data were Fourier transformed and CLEANed, using the Clark algorithm, and the restored Gaussian clean beam is $7''.9 \times 3''.8$, at a position angle of 14° . The results are shown in Figs. 1 and 2.

Images of the $\text{H}_2 \ v = 1 - 0 \ S(1)$ emission in the northern quadrant of the Helix nebula were obtained with the SOFI infrared camera on the ESO NTT, in June, 2001. The instrument has a 1024^2 Hawaii HgCdTe array and was used with an image scale of $0''.24 \text{ pixel}^{-1}$. The seeing was $1''.2$. The observations were made with an H_2 filter of width $0.028 \ \mu\text{m}$, centered at $2.124 \ \mu\text{m}$, well separated from the He I $2^1\text{P}_0-2^1\text{S} \ 2.058 \ \mu\text{m}$ and H I $2.166 \ \mu\text{m}$ lines, the latter having a transmission of a few percent in the filter passband. Fifteen 1 min exposures were made, using sky offset positions at $5'$ N, $5'$ E, and $7'$ NE. The individual images were sky cleaned, flat fielded, and combined with shift-and-add registration using the brightest stellar images.

Many globules are detected in the full H_2 image, and the region around globule C1 is shown in the top, center-left panel in Fig. 1. An approximate calibration has been made by comparing the smoothed, full image with the data of Speck et al. (2002). The intensity of the emission at the head of the globule is $\sim 10^{-4} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1}$. Astrometry of the H_2 and optical images was carried out using stars in the USNO catalog, and the registration between the images is $\sim 0''.1$ rms. The accuracy of the absolute positions for comparison with the CO interferometry is $\sim 0''.3-0''.5$.

3. Properties of the Globule

3.1. Overview

The observations presented in Fig. 1 provide complementary views of the Helix globule. The CO 1–0 line, with an upper level of $E_u = 5.5 \text{ K}$ above the ground state, shows the overall distribution and kinematics of the cool molecular gas; the high lying ($E_u \sim 7,000 \text{ K}$) $\text{H}_2 \ v = 1 - 0 \ S(1)$ line traces excited molecular gas; the image in [O III] shows the distribution of dust, seen in absorption against the nebular emission; and the image in $\text{H}\alpha + [\text{N II}]$ shows the photo-ionized surfaces facing the central star.

The observations immediately confirm the molecular nature of the globule, and reveal some important details of its structure.

3.2. CO Structure and Kinematics

The CO 1–0 velocity-integrated intensity map (Fig. 1) is seen to be extended with respect to the telescope beam, and the deconvolved source size, assuming a Gaussian distribution, is $\sim 2'' \times 10''$ ($1''$ corresponds to 3×10^{15} cm at 200 pc). The emission is marginally resolved in RA but is resolved in Dec, and extends roughly along the head-tail axis of the globule. The peak intensity is 2.6 K km s^{-1} , and the total flux is $166 \text{ K km s}^{-1} \text{ arcsec}^2$.

The CO velocity-strip (Fig. 2) shows that to the south, toward the head of the globule, the line width is extremely narrow, $\Delta V = 0.5 \text{ km s}^{-1}$ (FWHM), and it broadens out to $\sim 0.8 \text{ km s}^{-1}$ farther north into the tail region. The CO radial velocity is precisely determined to be $V_{\text{lsr}} = -27.9 \text{ km s}^{-1}$ (the correction to V_{hel} is -2.9 km s^{-1}), consistent with $-28.7 \pm 2 \text{ km s}^{-1}$ measured in $[\text{N II}] 6584\text{\AA}$ by Meaburn et al. (1998). The systemic velocity of the whole envelope of the Helix nebula is -23 km s^{-1} (Young et al. 1999), so the globule is blue-shifted by $\sim 5 \text{ km s}^{-1}$. For a radial expansion velocity of the globule system of $\sim 20 \text{ km s}^{-1}$ (Young et al. 1999), the globule lies on the near side of the nebula, on a radius vector from the star inclined to the line of sight by $\sim 75^\circ$. The head-tail axis, assumed radial, is also seen at this angle.

The CO channel maps (Fig. 1) show the structure in the molecular gas. There are two peaks, offset $3''$ N and $8''$ N from the field center. The first is just north of the maximum absorption in the $[\text{O III}]$ image (centered at $1''.5$ N), and we identify this molecular emission with the head of the globule whose photo-ionized surface toward the central star is seen in the $\text{H}\alpha + [\text{N II}]$ image. Substantial CO emission, however, extends away from the head into the tail region, and the second CO peak lies close to a second, weak maximum in the dust absorption image (at $8''$ N). This extended CO emission, together with the line broadening into the tail accounts for the offset of the peak in the CO integrated intensity map from the head of the globule.

These observations demonstrate that the molecular gas in the cometary globule is *not* a compact spheroidal bullet, but has a substantial component extending into the tail region.

3.3. H₂ Distribution and Excitation

The distribution of the H₂ emission in the globule is strikingly different from that of CO (see Fig. 1). The strongest H₂ emission occurs at the face of the globule toward the central star. There is little emission directly behind the globule, but observable emission extends to large distances ($\gtrsim 24''$) along the tail. Remarkably, the H₂ distribution most closely follows that of the ionized gas seen in $\text{H}\alpha + [\text{N II}]$.

The high spatial resolution of our observations and the well defined geometry of the globule provide a unique perspective on the question of the excitation of the H_2 (Cox et al. 1998), which affects its observed distribution. The bright H_2 emission clearly arises in a thin surface layer in the molecular gas. Fig. 3 shows a close-up of the globule head, comparing the H_2 and $\text{H}\alpha + [\text{N II}]$ emission with $[\text{O III}]$. The H_2 emission forms a limb-brightened cap on the globule facing the central star, and lies just inside ($0''.5\text{--}1''$) the halo of photo-ionized gas seen in $\text{H}\alpha + [\text{N II}]$. In the tail region too (Fig. 1), the H_2 is enhanced near two $\text{H}\alpha + [\text{N II}]$ peaks, at $8''$ N and $12''$ N on the east side, which are directly illuminated by the star. More detailed observations are needed to determine whether these are separate, small globules along the line of sight, or sub-structures of the main tail.

The observed distribution of the H_2 emission is in complete accord with the expectations of H_2 excitation in a thin photo-dissociation region (PDR) at the surface of the molecular gas. The observed intensity of the line is probably consistent with this interpretation (Natta & Hollenbach 1998; Speck et al. 2002) although detailed PDR models of the globules have not yet been developed. The possibility that the observed distribution is caused by shocks is unlikely, in view of the small crossing time ($\lesssim 400$ yr) for shocks ($v_s \gtrsim 5 \text{ km s}^{-1}$) able to excite the $v = 1 - 0$ $S(1)$ H_2 line (e.g., Burton et al. 1992); shocks in the bulk of the molecular gas are ruled out by the absence of any disturbance in the CO emission with velocities larger than $\sim 0.5 \text{ km s}^{-1}$.

3.4. Physical Properties of the Molecular Gas

Using the CO 2–1 observations of the globule by Huggins et al. (1992), we find the 2–1/1–0 flux ratio to be $\sim 2\text{--}3$ (assuming significant 1–0 flux is not resolved out on the shortest baselines), which suggests that the lines are at least partly optically thin, with an excitation temperature $\sim 18\text{--}40$ K in the thin limit. For a representative value of 25 K and a CO abundance of 3×10^{-4} , the CO 1–0 flux gives a mass of molecular gas in the globule of $M_{\text{m}} \gtrsim 1 \times 10^{-5} M_{\odot}$. The corresponding average density in a volume with projected dimensions of $2'' \times 10''$, which includes the head and the tail seen in CO, is $n_{\text{H}} \gtrsim 2 \times 10^4 \text{ cm}^{-3}$.

These values are consistent with a mass of $\sim 2 \times 10^{-5} M_{\odot}$ and $n_{\text{H}} \sim 4 \times 10^5 \text{ cm}^{-3}$ determined for the head of the globule by Meaburn et al. (1992) from the dust absorption seen in $[\text{O III}]$, corrected to a distance of 200 pc. The typical mass of photo-ionized gas at the surface of a globule, $\sim 10^{-9} M_{\odot}$ (e.g., O’Dell & Handron 1996), is negligible in comparison.

4. Origins and Evolution

The structure and kinematics of the molecular gas in the head and in the tail of the globule provide basic constraints on their origin and evolution.

One scenario for the origin of globules is that they form in the atmosphere of the progenitor star and are carried out in the expanding circumstellar envelope (Dyson et al. 1989). In this case, the wind-swept appearance of the globules suggests a model in which material from the head is swept into the tail by a radially directed wind, and Dyson, Hartquist, & Biro (1993) have shown that the wind needs to be subsonic to form a narrow tail. Our observations constrain this model in showing no evidence for a wind-swept flow pattern at the present time. There is no difference in velocity between the CO emission in the head and the tail of the globule, and from the CO strip map (Fig. 2) the differential motion is $\lesssim 0.2 \text{ km s}^{-1}$ along the line of sight, or $\lesssim 0.8 \text{ km s}^{-1}$ in a radial direction, corrected for the inclination. This would produce a tail of length $\lesssim 4''$ in 5,000 yr which is likely the maximum time available for such a process. The tails could have been fully formed at an earlier epoch, or it might be that ablation occurs only from the ionized edges of the head of the globule, but this would not account for the presence of molecular gas seen in H_2 along the whole length of the tail.

A different view of the gas dynamics around the globule at the present time is provided by the photo-evaporation model, which has been discussed in the context of the Helix globules by López-Martín et al. (2001). In this model, photo-ionization of the neutral globule, whose molecular core is unambiguously confirmed by our observations, produces an ionized outflow from the surface. Given that the density of the ionized gas at the head of the globule is $\sim 10^3 \text{ cm}^{-3}$, and that of the ambient gas is much lower $\sim 50 \text{ cm}^{-3}$ (e.g., O’Dell & Handron 1996), any subsonic flow of the ambient gas around the globule is unlikely to have important dynamical effects in shaping the gas in the tail.

An alternative mechanism for growing a tail on a pre-existing globule is by shadowing (Cantó et al. 1998), which can form a tail behind the globule as the result of under-pressure of the gas in the shadowed region. For the simplest case of a neutral globule in ionized gas, this model produces only modest density increases in the tail, and is unlikely to lead to the substantial molecular tails revealed by the present observations. Striking effects of shadowing of the ionizing radiation are, however, seen in optical images of the Helix (e.g., Henry, Kwitter, & Dufour 1999), where long, radial plumes appear as extensions to globules and their tails. Similar shadowing of the stellar radiation at wavelengths longer than the Lyman limit must also occur, and the consequent reduction of the CO and H_2 photo-dissociation rates in the shadowed regions must play a key role in preserving the molecular gas in the extended tails.

If globules originate close to the star as proposed by Dyson et al. (1989), the most crucial phase in their evolution occurs when they are overrun by the ionization front of the nebula. At this stage, the shadowing of pre-existing molecules or hydrodynamic flows, or both, could lead to globules with molecular tails. An alternative scenario is one in which the globules and their tails originate simultaneously in instabilities at the ionization front (e.g., Capriotti 1973). In this model, shadowing and possibly hydrodynamic flows would also likely play a role, and the fingers of gas which result from the instability could contribute directly to the formation of the tails. The later development of these two models will likely be similar, and can plausibly lead to the globules with substantial molecular tails that we now observe. Further studies should allow us to discriminate between them.

As for the fate of the globules, Meaburn et al. (1998) estimated an ablation rate for globule C1 of $3 \times 10^{-8} \text{ M}_{\odot} \text{ yr}^{-1}$ based on densities inferred from the [O III] image, and an assumed ablation flow velocity of 10 km s^{-1} . This implies an improbably short lifetime of $\sim 500 \text{ yr}$ which led Meaburn et al. to consider ad hoc, restricted flow models around the globule. The absence of an observable ablation flow in the globule reported here for CO ($\lesssim 0.8 \text{ km s}^{-1}$) completely resolves this problem, by increasing the original estimate of the ablation time scale by at least an order of magnitude. In fact, the current mass-loss rate of the globule is likely dominated by photo-evaporation (López-Martín et al. 2001), and the corresponding time scale is $\sim 10^4 \text{ yr}$. Thus the globules are long lived, and may even escape the nebula.

We thank the staff of the Plateau de Bure interferometer for making the observations. This work has been supported in part by NSF grant AST-9986159 (to P.J.H.).

REFERENCES

- Burton, M. G., Hollenbach, D. J., & Tielens, A. G. G. M. 1992, *ApJ*, 399, 563
- Cantó, J., Raga, A., Steffen, W., & Shapiro, P. R. 1998, *ApJ*, 502, 695
- Capriotti, E.R. 1973, *ApJ*, 179, 495
- Cox, P., et al. 1998, *ApJ*, 495, L23
- Dyson, J. E., Hartquist, T. W., Pettini, M., & Smith, L. J. 1989, *MNRAS*, 241, 625
- Dyson, J. E., Hartquist, T. W., & Biro, S. 1993, *MNRAS*, 216, 430
- Forveille T., & Huggins P. J. 1991, *A&A*, 248, 599
- Harris, H. C., Dahn, C. C., Monet, D. G., & Pier, J. R. 1997, in *IAU Symp. 180, Planetary Nebulae*, ed. H. J. Habing & H. J. G. L. M. Lamers (Dordrecht: Kluwer), 40

- Henry, R. B. C., Kwitter, K. B., & Dufour, R. J. 1999, *ApJ*, 517, 782
- Huggins, P. J., Bachiller, R., Cox, P., & Forveille, T. 1992, *ApJ*, 401, L43
- Kastner, J. H., Weintraub, D. A., Gatley, I., Merrill, K. M., & Probst, R. G. 1996, *ApJ*, 462, 777
- López-Martín, L., Raga, A. C., Mellema, G., Henney, W. J., & Cantó, J. 2001, *ApJ*, 548, 288
- Meaburn, J., Walsh, J. R., Clegg, R. E. S., Walton, N. A., Taylor, D., & Berry, D. S. 1992, *MNRAS*, 255, 177
- Meaburn, J., Clayton, C. A., Bryce, M., Walsh, J. R., Holloway, A. J., & Steffen, W. 1998, *MNRAS*, 294, 201
- Natta, A., & Hollenbach, D. 1998, *A&A*, 337, 517
- O’Dell, C. R., & Handron, K. D. 1996, *AJ*, 111, 1630
- O’Dell, C. R., Balick, B., Hajian, A. R., Henney, W. J., & Burkert, A. 2002, *AJ*, in press
- Rodriguez, L. F., Goss, W. M., & Williams, R. 2002, *ApJ*, in press
- Speck, A. K., Meixner, M., Fong, D., McCullough, P. R., Moser, D. E., & Ueta, T. 2002, *AJ*, 123, 346
- Walsh, J. R., & Meaburn, J. 1993, *The Messenger*, 73, 35
- Young, K., Cox, P., Huggins, P. J., Forveille, T., & Bachiller, R. 1999, *ApJ*, 522, 387

Fig. 1.— Observations of the Helix globule C1. Top row: Images in CO 1–0 integrated intensity (left); H_2 $v = 1 - 0$ $S(1)$ (center left); $\text{H}\alpha + [\text{N II}]$ 6584 Å (center right); and dust absorption, seen against the nebula emission in $[\text{O III}]$ 5007 Å (right). Bottom row: CO 1–0 channel maps; the channels are 0.2 km s^{-1} wide and are centered at the velocities given in the upper right of each panel. The contour intervals are 0.4 K km s^{-1} for the CO integrated intensity map, and 0.9 K for the channel maps. The ellipse in the top, left panel shows the beam size for the CO observations. For each panel, the offsets are relative to a field center of 22:29:37.78 –20:48:02.00 (J2000) which is used for all images in the paper.

Fig. 2.— CO 1–0 velocity strip map of the globule along the major axis. The contour intervals are 0.5 K .

Fig. 3.— Close-up of the head of the globule. Contours of the emission in H_2 $v = 1 - 0$ $S(1)$ (left) and $\text{H}\alpha + [\text{N II}]$ (right), superposed on the $[\text{O III}]$ image. The H_2 data are smoothed with a Gaussian $0''.6$ (FWHM). The contours are 0.3, 0.5, 0.7, and 0.9 of the peak emission.





